

Remplir[™]

DESCRIPTION

Remplir[™] is a sterile, resorbable collagen membrane intended for use in peripheral nerve repair.

Remplir[™] is a purified collagen membrane of porcine origin produced using quality-controlled manufacturing processes. Raw materials are sourced exclusively from within Australia and collected at certified facilities under strict quality controls. The collagen is purified to remove materials that may elicit antigenic reactions. Remplir[™] is manufactured without crosslinking. **INDICATIONS FOR USE**

Remplir[™] is indicated for the repair of peripheral nerve injuries.

PROPERTIES

Remplir[™] is designed to protect the injured nerve following repair and provide a favourable environment for nerve regeneration to occur. Remplir[™] permits passage of fluids but acts as a barrier to cellular ingrowth.

Remplir[™] is soft and pliable with a smooth outer surface and rough inner surface, providing excellent handling characteristics. Remplir[™] is wettable and may be sutured in place, if required.

DIRECTIONS FOR USE

These directions are intended as a general guide and are not intended to supersede professional clinical judgement.

- 1. Follow standard procedures for nerve exposure and mobilization. Perform neurolysis and/or coaptation of nerve ends if required.
- 2. Trim Remplir[™] to required size.
- 3. Hydrate if required with sterile saline or other physiological solution.
- 4. Wrap Remplir[™] around the site of the nerve injury/repair without constricting or compressing the nerve.
- 5. If necessary, fix Remplir[™] in place with atraumatic sutures.
- 6. Follow standard post-operative procedures and care. Application of Remplir[™] does not modify post-operative treatment.

LIMITATIONS OF USE

CONTRAINDICATIONS

The use of Remplir[™] in patients with known sensitivity to porcine-derived materials or collagen is contraindicated.

ADVERSE REACTIONS

Possible complications of peripheral nerve surgery include pain, infection, swelling, inflammation, foreign body sensation and changes in nerve sensitivity (paraesthesia or hyperaesthesia) at the implant site.

Adverse reactions to porcine-derived collagen nerve wraps such as Remplir[™] are extremely rare. Patients should be monitored closely in the initial post-operative period to identify and address any adverse reactions that may occur.

PRECAUTIONS

Safety and efficacy for use in patients <18 years cannot be assured.

Peripheral nerve repair with Remplir[™] should only be undertaken by suitably qualified surgeons.

STORAGE AND HANDLING

Store Remplir[™] in its original packaging at room temperature (15-25°C) in a dry place.

HOW SUPPLIED

Remplir[™] is supplied in a double PETG/Tyvek blister pack contained in a labelled cardboard box.

Size	Code
15x20mm	ON-152
20x30mm	ON-203
30x40mm	ON-304
40x50mm	ON-405

MANUFACTURER



Orthocell Ltd Building 191, Murdoch University South Street, Murdoch WA 6150 Australia

HKMD No. 251574 没有中文版使用説明

Asia Actual Limited Local Responsible Person 17/F, Chung Pont Commercial Bldg, 300 Hennessy Road, HK PH: +852 2893 3118

Asia Actual Limited 本地負責人 灣仔軒尼詩道300號 中邦商業大厦17樓 聯絡電話: +852 2893 3118

SYMBOLS USED IN LABELLING

52 Use-by date

STEPSUZE

LOT

(2)



REF Catalogue number

w Date of manufacture

STERILE R Sterile using irradiation

Double sterile barrier system

> 鯊 Keep away from sunlight



Do not resterilize

Temperature limit (15-25°)



Do not re-use



is damaged and

Do not use if package

\$

Medical Device



Contains biological

material of animal

BIO

origin

Identifier





T

Keep dry

